



Development of Endosomal Escape Vehicles to Enhance the Intracellular Delivery of Oligonucleotides

Patrick Dougherty, PhD
Senior Scientist, Discovery Research

Boulder Peptide Symposium
November 10th, 2022



This presentation includes express and implied “forward-looking statements. Forward looking statements include all statements that are not historical facts, and in some cases, can be identified by terms such as “may,” “might,” “will,” “could,” “would,” “should,” “expect,” “intend,” “plan,” “objective,” “anticipate,” “believe,” “estimate,” “predict,” “potential,” “continue,” “ongoing,” or the negative of these terms, or other comparable terminology intended to identify statements about the future. Forward-looking statements contained in this presentation include, but are not limited to, statements about our product development activities and clinical trials, our regulatory filings and approvals, our ability to develop and advance our current and future product candidates and discovery programs, our ability to establish and maintain collaborations or strategic relationships or obtain additional funding, the rate and degree of market acceptance and clinical utility of our product candidates, the ability and willingness of our third-party collaborators to continue research and development activities relating to our product candidates, our and our collaborators’ ability to protect our intellectual property for our products. By their nature, these statements are subject to numerous risks and uncertainties, including factors beyond our control, that could cause actual results, performance or achievement to differ materially and adversely from those anticipated or implied in the statements. You should not rely upon forward-looking statements as predictions of future events. Although our management believes that the expectations reflected in our statements are reasonable, we cannot guarantee that the future results, performance or events and circumstances described in the forward-looking statements will be achieved or occur. Recipients are cautioned not to place undue reliance on these forward-looking statements, which speak only as of the date such statements are made and should not be construed as statements of fact.


Certain information contained in this presentation and statements made orally during this presentation relate to or are based on studies, publications, surveys and other data obtained from third-party sources and our own internal estimates and research. While we believe these third-party studies, publications, surveys and other data to be reliable as of the date of this presentation, it has not independently verified, and makes no representations as to the adequacy, fairness, accuracy or completeness of, any information obtained from third-party sources. In addition, no independent source has evaluated the reasonableness or accuracy of our internal estimates or research and no reliance should be made on any information or statements made in this presentation relating to or based on such internal estimates and research.

ENTRADA'S MISSION


*Treating Devastating Diseases With
Intracellular Therapeutics*

THE NEED FOR INTRACELLULAR THERAPEUTICS

Approximately 75% of all disease-causing targets are intracellular and difficult to reach, representing a significant issue when developing effective therapies




First Challenge:
Cell Entry



**The Endosomal Escape Vehicle (EEV™) Platform aims to solve the fundamental problem:
Lack of efficient cellular uptake and escape from the endosome**

Entrada's EEV platform consists of a library of proprietary cyclic peptides with unique chemistry that enable improved uptake and endosomal escape

- Unique chemistry results in **improved uptake and endosomal escape**
- Cyclic structure enhances **proteolytic stability**
- Small and cyclic structure may **reduce immunogenicity risk**
- Mechanism of internalization **conserved across species**




A BROADLY APPLICABLE PLATFORM

Entrada has demonstrated intracellular uptake of unique moieties ranging from 1 kDa to 600 kDa


Antibodies



Enzymes




Oligonucleotides




550-600 KDa

Hybrid frataxin



150 KDa

Antibody




98 KDa

Thymidine
phosphorylase




96 KDa

Purine
nucleoside
phosphorylase




86 KDa

Alanine-
glyoxylate
aminotransferase




46 KDa

Human frataxin




37 KDa

PTP1B
catalytic
domain



32 KDa

EGFP



16 KDa

Nanobody



6 KDa


Oligonucleotide



1-3 KDa


Various
peptide cargos

Discovery Engine for Intracellular Therapeutics




- Cyclic peptide library design and combinatorial synthesis to generate **EEV library**
- Delivery and counter-screening assays enabled for *in vitro* **high throughput screening**
- Functional screening of lead EEVs *in vivo* to select for **pharmacodynamic activity** in target tissues
- Optimize **linker & conjugation chemistry** for desired therapeutic modality

Screening Cascade for EEV Candidates




Entrada's EEV-therapeutics can be designed to enhance functional delivery to target tissues

EGFP-654 Transgenic Mice



Functional Delivery to Target Tissues



TRANSLATION FROM UPTAKE TO OUTCOMES

EEV-therapeutics possess favorable pharmacological properties: significant uptake in target tissues, efficient intracellular delivery and potent pharmacodynamic outcomes

Tissue Uptake in Muscle



- ✓ Skeletal muscle
- ✓ Cardiac muscle

Intracellular Delivery




- ✓ Endosomal escape
- ✓ Nuclear localization


Pharmacodynamic Outcome




- ✓ Rapid, dose-dependent efficacy
- ✓ Duration of at least 12 weeks



IV, hDMD mice, 5-day p.i.



24-hour incubation




IV, hDMD mice, 5-day p.i.

hDMD mice express full-length human dystrophin gene. p.i. post injection; shown as mean ± standard deviation.

DUCHENNE MUSCULAR DYSTROPHY (DMD)

Duchenne muscular dystrophy (DMD) is a progressive, devastating muscle wasting disease with significant unmet need. Entrada's first DMD program is for exon 44 skipping


- DMD is caused by mutations in the *DMD* gene that encodes for dystrophin¹
- Progressive muscle degeneration, wasting and paralysis generally lead to death via **respiratory and/or cardiac failure**²
- Exon skipping therapeutics (for exons 45, 51 and 53) using **PMO chemistry** were approved based on **a very modest improvement in dystrophin levels ranging from ~1 to 6%**
- 40% of patients with DMD have mutations amenable to exon skipping of exons 44, 45, 51 and 53.³ **However, there is currently no approved therapy for patients with mutations amenable to exon 44 skipping**




REPEAT EEV-PMO TREATMENT IN D2-*mdx* MICE

Robust exon 23 skipping after four monthly IV doses of EEV-PMO-23 in D2-*mdx* mice


Broad dystrophin expression and restoration of muscle integrity after four monthly IV doses of EEV-PMO-23 in D2-*mdx* mice



Representative Immunofluorescence of Gastrocnemius Muscle (Dystrophin Staining in Green)



Representative Histopathology of Gastrocnemius Muscle (H&E Staining)




- D2-*mdx* mice (male, n=6-7) were treated with 4 monthly doses of either vehicle, 20 mg/kg PMO-23 or 20 mg/kg PMO-23 equivalent of EEV-PMO-23, and the data were collected ~4 weeks after the last dose


REPEAT EEV-PMO TREATMENT RESULTS IN CK CORRECTION AND FUNCTIONAL IMPROVEMENT

Repeat EEV-PMO-23 treatment normalized serum CK levels and showed significant improvements in muscle function when compared to PMO alone after four monthly IV doses in D2-*mdx* mice


Serum CK Levels



Wire Hang Time



Grip Strength




- **D2-*mdx*** is a DMD mouse model on DBA/2J background and better recapitulate disease pathology (Fukada et al. *Am. J. Path.* 2010)
- D2-*mdx* mice (male, n=6-7) were treated with 4 monthly doses of either vehicle, 20 mg/kg PMO-23 or 20 mg/kg PMO-23 equivalent of EEV-PMO-23, and the data were collected ~2 weeks after the last dose

CK, creatine kinase; *p<0.05, **p<0.01, ****p<0.0001; n.s., not significant; shown as mean ± standard deviation.


EXON SKIPPING AND DYSTROPHIN PRODUCTION IN PATIENT-DERIVED CELLS

Robust dose-dependent exon 44 skipping and dystrophin protein production were observed in DMD patient-derived muscle cells treated with clinical candidate, ENTR-601-44

Exon Skipping




Dystrophin Protein Production



DMDΔ45, immortalized myoblasts from DMD patients harboring an out-of-frame exon 45 deletion and further differentiated into myotubes; ENTR-601-44 is the clinical candidate selected for DMD exon 44 skipping EEV-oligonucleotide conjugate; ****p<0.0001; shown as mean ± standard deviation.

ENTR-601-44 IN HUMAN DMD MICE (hDMD)


A single IV dose of ENTR-601-44 resulted in high and durable levels of exon 44 skipping across major muscle groups in hDMD mice for at least 12 weeks



- Post single IV dosage of 60 mg/kg (PMO equivalent), robust exon skipping was observed in ENTR-601-44 treated hDMD mice


A single IV dose of ENTR-601-44 resulted in robust exon skipping in both skeletal and cardiac muscles in NHP, as well as prolonged duration of effect for at least 12 weeks

Exon Skipping in NHP Muscles at Day 7



- At 7 days post IV infusion of 30 mg/kg (PMO equivalent), robust exon 44 skipping across different muscle groups isolated from the ENTR-601-44 treated NHP

Duration of Effect in NHP Biceps for at Least 12 Weeks



- Post IV infusion of 35 mg/kg (PMO equivalent), robust exon 44 skipping observed in biceps in the ENTR-601-44 treated NHP (n=3 per cohort) for at least 12 weeks

NHP, non-human primates; shown as mean \pm standard deviation.


Entrada's murine and NHP data represent a robust set of translational data;
IND application in Q4 2022

- High levels of exon skipping across *mdx*, *D2-mdx*, human dystrophin mouse and NHP studies
- Exon skipping translates to promising dystrophin production in heart and skeletal muscles
- Dystrophin production sufficient to result in functional improvement
- Extended circulating half-life and durable dystrophin production over 12+ weeks from a single injection of ENTR-601-44 was observed in the NHP
- ENTR-601-44 IND application planned in Q4 2022
- Exon 45 candidate expected to enable second DMD candidate selection in quick succession

MYOTONIC DYSTROPHY TYPE 1 (DM1)

DM1 is a debilitating multi-systemic disease with no available treatments; CUG repeats in DMPK mRNA sequester MBNL proteins, resulting in nuclear foci, aberrant splicing, and disease


- **DM1 occurs in 1:8,000 people worldwide and affects ~40,000 patients in US and over 50,000 in Europe**
 - 75% of patients have adult-onset DM1
 - Multisystemic: including myotonia, muscle weakness and atrophy, cardiac conduction abnormalities, pulmonary complications, cataracts, and endocrine dysfunction¹
 - **Currently there are no approved therapies**
- **DM1 is caused by CUG repeats in the mRNA that sequester MBNL proteins²**
 - Mutant *DMPK* mRNA and MBNL proteins form aggregates named nuclear foci³
- **MBNL activity is decreased as a result of sequestration leading to spliceopathy of downstream transcripts^{4,5}**




ENTR-701 IN DM1 CELL LINE WITH REPEAT EXPANSION

DM1 clinical candidate, ENTR-701, showed reduction of nuclear foci and selective reduction of repeat expansion-containing DMPK transcript in the HeLa480 cell line

Nuclear Foci Reduction



Selective Reduction of Mutant DMPK mRNA




- Free uptake of ENTR-701 reduced nuclear foci and selectively reduced (CUG)480 containing DMPK mRNA



ENTR-701 treatment in DM1 patient-derived muscle cells resulted in significant nuclear foci reduction and correction of aberrant splicing

Nuclear Foci Reduction

DM1 cells untreated




DM1 cells ENTR-701




Correction of Aberrant Splicing

MBNL1




SOS1



- Immortalized DM1 patient-derived (2,600 CUG repeats) muscle cells¹ were treated with ENTR-701 and analyzed for the reduction of nuclear foci and the correction of aberrant splicing


¹Arandel, L. et al. *Dis. Model. Mech.* 2017. **ENTR-701** is the clinical candidate selected for DM1, composed of CUG-repeat blocking PMO conjugated to EEV; ****p<0.0001 for ENTR-701 compared to untreated; shown as mean ± standard deviation.

ENTR-701 treatment reduced nuclear foci and CUG-repeat expansion transcript level and corrected aberrant splicing in HSA-LR mice



- HSA-LR mice were dosed with vehicle or ENTR-701 and taken down 1-week post injection; tibialis anterior samples analyzed as a representative skeletal group

ENTR-701 CORRECTED SPLICEOPATHY IN HSA-LR MICE



DM1-affected splicing events analyzed by RNA-seq; **ENTR-701** is the clinical candidate selected for DM1, composed of CUG-repeat blocking PMO conjugated to EEV




DURABILITY OF ENTR-701 IN HSA-LR MICE

A single dose of ENTR-701 resulted in splicing correction in HSA-LR mice for at least 8 weeks

Mbn1 Exon 5 Inclusion

Atp2a1 Exon 22 Inclusion

Nfix Exon 7 Inclusion



■ Wild type ■ HSA-LR + Vehicle ■ HSA-LR + ENTR-701

- Post single IV dosage of 60 mg/kg (PMO equivalent), robust splicing correction in the ENTR-701 treated HSA-LR mice 4, 6, or 8 weeks post injection

ENTR-701 is the clinical candidate selected for DM1, composed of CUG-repeat blocking PMO conjugated to EEV; *Mbn1*, muscleblind like splicing regulator 1; *Atp2a1*, sarcoplasmic/endoplasmic reticulum calcium ATPase; *Nfix*, nuclear factor I X; ***p<0.001, ****p<0.0001, shown as mean ± standard deviation.

MYOTONIA CORRECTION IN HSA-LR MICE

A single dose of ENTR-701 ameliorated pinch-induced myotonia for at least 8 weeks

HSA-LR Mouse: **Non-treated**



HSA-LR Mouse: **ENTR-701 Treated**









ENTR-701 demonstrated potential to treat DM1 via a CUG-repeat steric blocking approach both *in vitro* and *in vivo*; IND application planned in 2023

- Robust *in vitro* and *in vivo* data set demonstrating:
 - Highly specific reduction of pathogenic CUG-repeat containing mRNA
 - Reduction of nuclear foci
 - Correction of *Mbn1* and downstream aberrant splicing
 - Correction of global transcriptome
- Single dose of ENTR-701 demonstrates durable splicing correction and amelioration of myotonia for at least 8 weeks post-dose in HSA-LR model
 - Importantly, HSA-LR model carries a transgene resulting in CUG repeat expansion and recapitulates DM1 phenotype and molecular pathology
- ENTR-701 IND submission planned in 2023

DISCOVERY PROGRAMS

ADDITIONAL PLATFORM OPPORTUNITIES

Entrada continues to invest in and build upon our EEV Platform to extend our efforts in developing novel EEV-therapeutic candidates

Target		Platform Approach		Goal
	DNA		Gene editing	Deliver CRISPR enzyme and repair gene function with guide RNA
	RNA		RNA editing	Deliver oligonucleotide therapeutics for RNA editing
			RNA splicing	Modify RNA via exon/intron splicing to activate protein expression
			RNA blocking	Block trinucleotide repeats in RNA to inhibit adverse binding
			RNA silencing	Silence or knockdown RNA to prevent protein expression
	Protein		Protein replacement	Replace proteins and enzymes
			Protein inhibition	Inhibit protein signaling pathways
			Protein degradation	Degrade disease-causing proteins

Thank you!



WANT TO LEARN MORE?

Scan the QR code for access to our recent publications and conference presentations.

